

Chemistry—A European Journal

Supporting Information

An α 2,3-Sialyltransferase from *Photobacterium phosphoreum* with Broad Substrate Scope: Controlling Hydrolytic Activity by Directed Evolution

Alexander Mertsch, Ning He, Dong Yi, Michael Kickstein, and Wolf-Dieter Fessner*^[a]

Supporting Information

Index	page
Cloning, expression, and purification of 2,3SiaT_{pph}	2
Figure S1. Plasmid construction for expression of 2,3SiaT _{pph}	2
Figure S2. Cloning and expression of 2,3SiaT from <i>Photobacterium phosphoreum</i>	2
Table S1: Primers for site directed mutagenesis library	3
Table S2. Primers for second generation combination variants	4
Table S3. Templates and primer mix combination for second generation combination library	4
Table S4. Primers for preparation of specific individual combination variants	4
Table S5. Primer and template combination for specific individual combination variants	4
Table S6. Analysis of distances between active site residues and substrates or water molecules	5
Table S7. Thermodynamic stability of 2,3SiaT _{pph} determined by nanoDSF	5
HPTLC screening	6
Figure S3. Elution by <i>n</i> -propanol:H ₂ O:NH ₃ = 7:3:1 for unbiased integration of Neu5Ac	6
Figure S4. Quantitative analysis of sample from Figure S3; inset shows integration of lane 2	6
Figure S5. Elution by <i>n</i> -butanol:acetone:H ₂ O:AcOH = 35:35:23:7 for unbiased integration of SiaLac	7
Figure S6. Quantitative analysis for lane 1	7
Table S8. HPTLC analysis of sialyltransfer vs. hydrolysis using 2,3SiaT _{pph} and variants (duplicate runs)	8
Control for CMP independent sialidase activity	
Table S9. Data for Figure 4: Quantification of sialidase activity	9
Figure S7. Control experiment for sialidase activity in the presence of alkaline phosphatase	9
General protocol for synthesis of fluorescence-labeled neo-sialoconjugates using native 2,3SiaT_{pph}	10
Synthesis of α2,3-SiaLac (GM3 trisaccharide, 14) by engineered 2,3SiaT_{pph} (A151D)	17
Exemplary kinetic profiles for native enzyme and best variants	19
Quality control of second-generation combination library by sequencing	20

Cloning, expression, and purification of 2,3SiaT_{pph}




Figure S1. Plasmid construction for expression of 2,3SiaT_{pph}.




Figure S2. Cloning and expression of 2,3SiaT from *Photobacterium phosphoreum*. a) Expression level of the recombinant SiaT_{pph} in BL21(DE3) analyzed by SDS-PAGE. Lanes 1: marker; 2: crude cell extract; 3: supernatant of crude cell extract. b) Purification of 2,3SiaT_{pph} by Ni-affinity chromatography. Lanes 1: marker; 2: purified 2,3SiaT_{pph}.

Table S1. Primers for site directed mutagenesis library (Agilent method)

SiaT-L120NNK_F	AGTCATTATTAACGGCAATACGnnkTGGCTGTCGACGTGGTAAAC
SiaT-L120NNK_R	GTTTACCACGTCGACAGCCCCmnnCGTATTGCCGTTAATAATGACT
SiaT-W121NNK_F	GTCATTATTAACGGCAATACGCTGnnkGCTGTCGACGTGGTAAACATTATC
SiaT-W121NNK_R	GATAATGTTTACCACGTCGACAGCmnnCAGCGTATTGCCGTTAATAATGAC
SiaT-D148NNK_F	AACCGAGATCGAACTGAACTTCTATGATnnkGGTAGCGCCGAGTAT
SiaT-D148NNK_R	ATACTCGGCCTACCmnnATCATAGAAGTTCAGTCGATCTCGTT
SiaT-S150WVC_F	CGAACTGAACCTCTATGATGATGGTwvcGCCGAGTATGTT
SiaT-S150WVC_R	GAACATACTCGGCGbwaCCATCATCATAGAAGTTCAGTCG
SiaT-A151NNK_F	CTGAACTTCTATGATGATGGTAGCnnkGAGTATGTTGCCTGTATGATTTC
SiaT-A151NNK_R	GAAATCATACAGGCGAACATACTCmnnGCTACCATCATAGAAGTTCAG
SiaT-R155NNK_F	GATGATGGTAGCGCCGAGTATGTTnnkCTGTATGATTTCTCTCGC
SiaT-R155NNK_R	GCGAGAGAAATCATACAGmnnAACATACTCGGCGCTACCATCATC
SiaT-N190NNK_F	CATCAACGGTACTCAGCCGTTGATnnkAGCATTGAAAACATCTATGG
SiaT-N190NNK_R	CCATAGATGTTTCAATGCTmnnATCGAACGGCTGAGTACCGTTGATG
SiaT-T278WVC_F, SiaT-N279WVC_F	CACGAAAACTTCATCTTCATTGGCwvcwvcTCTGGTACTGCCACTGCCGAGCAA
SiaT-T278WVC_R, SiaT-N279WVC_R	TTGCTCGCAGTGGCAGTACCAAGAGbwgbwgCCAATGAAGATGAAGTTTCGTG
SiaT-H317NNK_F	TGGCACCAATTCTGGTACTGCnnkTGCCGAGCAACAAATCGATAT
SiaT-H317NNK_R	ATATCGATTGTTGCTCGGCAmnnGCAGTACCAAGAATTGGTGCCA
SiaT-S359 WVC, S360WVC_F	CAGATGCCGTGGGTGGTATGGGTwvcwvcCGTTTCTTCACTGCCAAAGAC
SiaT-S359 WVC, S360WVC_R	GTGTTGGCAGTGAGAAGAAAACGbwgbwgACCCATACCACCCACGGCATCTG
SiaT-L364NNK_F	ACCGACATTGAGAATAACGCCnnkATCCAGGTATGAGCTG
SiaT-L364NNK_R	CAGCTCAATCATGACCTGGATmnnGGCGTTATTCTCAATGTCGGT

Table S2. Primers for second generation combination variants (NEB method)

1) SiaT-150S/T, SiaT151L/M_F	ascmtgGAGTATGTTGCCGTATG
2) SiaT-D148S/L_R	accraATCATAGAAGTTCAGTCG
3) SiaT-S150S/T, A151_F	ascgccGAGTATGTTGCCGTATG
4) D148_R	accatCATCATAGAAGTTCAGTCG

Primer mix: A = 1+2, B = 1+4, C = 2+3, D = 3+4

Table S3. Templates and primer mix combination for second generation combination library

Primer mix → Position ↓	A	B	C	D	template: native gene
	A	B	C	D	
D148	L/S	D	L/S	D	template: S359T/S360T
S150	S/T	S/T	S/T	S/T	
A151	L/M	L/M	A	A	

Table S4. Primers for preparation of specific individual combination variants (NEB method)

SiaT-A151D_F	TGATGGTAGC gat GAGTATGTTCG
SiaT-A151D_R	TCATAGAACGTTCAGTCGATCTCGG
SiaT-E342A_F	AATCCCGTTC gcg GCCCTGATTATG
SiaT-E342A_R	TTGTTATAAAATCTCGATCATGTTATG
SiaT-S359T_F	TGGTATGGGT acc AGCGTTTCTTCAC
SiaT-S359T_R	CCCACGGCATCTGGCAGAG
SiaT-L387A_F	GAATAACGCC gcg ATCCAGG
SiaT-L387A_R	TCAATGTCGGTTCGCTTTATAG

Table S5. Primer and template combination for specific individual combination variants

template	primer	product
w.t.	SiaT-E342A_F/R	E342A
A151D	SiaT-E342A_F/R	A151D/E342A
A151D	SiaT-L387A_F/R	A151D/L387A
S359T/S360T	SiaT-A151D_F/R	A151D/S359T/S360T
A151D/S359T/S360T	SiaT-L387A_F/R	A151D/S359T/S360T/L387A

Table S6. Analysis of distances between 13 selected amino acid residues and substrates or water molecules in the active site, and distances between water molecules and the centers of nucleophile attack during transfer or potential hydrolysis (sialoside C2 and phosphate, respectively).

AA position / Codon type	Contact distances [Å] ^{a)}						
L120 / NNK	CD2, GalO2: 4.7	CD2, GalO3: 5.0	CD2, GalO4: 4.1				
W121 / NNK	NE1, GalO4: 3.4	NE1, GalO5: 3.7	NE1, GalO6: 3.6				
D148 / NNK	OD2, GalO3: 2.1	OD2, GalO4: 2.4	OD2, SiaO4: 3.4	OD2, wat4: 4.5			
	OD1, GalO3: 3.7	OD1, GalO4: 1.9	O, SiaO4: 4.0	O, wat4: 2.7			
S150 / WVC	OG, SiaO1: 4.8	OG, SiaO4: 3.5	OG, PO2: 4.2	OG, wat1: 2.8	OG, wat2: 4.6		
	N, SiaO1: 4.3	N, SiaO4: 2.1	OG, SiaN5: 1.7				
A151 / NNK	N, SiaO10: 3.9	N, SiaN5: 3.0	N, SiaN5: 3.0		CB, GalO3: 4.3		
R155 / NNK	NH2, GalO1: 6.5	NE, GalO2: 7.8	NE, SiaO10: 7.3	NH2, GluO1: 5.1			
N190 / NNK	ND2, GalO1: 7.2	ND2, GalO2: 9.1	ND2, GluO1: 6.4				
T278 / WVC	OG1, wat3: 2.8	OG1, wat2: 6.1	OG1, SiaO9: 4.7	OG1, SiaO8: 5.5			
N279 / WVC	N, SiaO9: 6.6	O, SiaO9: 7.0	ND2, SiaO9: 5.8	OD1, SiaO9: 5.5			
H317 / NNK	NE2, PO3: 4.1	NE2, PO4: 6.3	NE2, PA: 5.6	NE2, SiaO8: 4.3	NE2, SiaO9: 5.4	N, CMPN4: 4.7	
S359 / WVC	OG, PO2: 3.5	OG, PO3: 3.7	OG, PO4: 3.7	OG, PO4: 3.9	OG, wat3: 4.0		
					N, wat3: 4.3	N, wat2: 4.8	
S360 / WVC	N, PO2: 1.3	N, PO3: 3.7	N, PA: 2.7	N, PO4: 3.4	N, wat1: 3.6	N, wat2: 5.6	
	OG, PO2: 1.8	OG, PO3: 3.4	OG, PA: 2.2	OG, PO4: 3.2	OG, wat1: 3.8	OG, wat4: 5.3	
	OG, PO1: 1.8						
L387 / NNK	CD2, wat1: 4.7	CD2, wat2: 4.0		CD1, SiaC11: 4.1	CD2, SiaC11: 5.2	CD2, PA: 8.7	
	Distances between water molecules and phosphate/SiaC2						
	PA, wat1: 4.4	PA, wat2: 6.4	PA, wat3: 5.7	PA, wat4: 6.2			
	SiaC2A, wat1: 5.1	SiaC2A, wat2: 6.9	SiaC2A, wat3: 7.2	SiaC2A, wat4: 4.8			

[a] Based on the 3D alignment model, as determined by PyMOL analysis. Selections contain: CD2 (contact in AA), GalO2 (contact to substrate and water); C: contact atom underlined, CD2: PyMOL atom descriptor. For clarity, PyMOL descriptors were amended by prefix Sia for Neu5Ac moiety, Gal for galactosyl unit and Glu for glucose unit in lactose.

Table S7. Thermodynamic stability of 2,3SiaT_{pph} determined by nanoDSF.

enzyme variant	melting point T_m [°C]
parent	47.1
A151D	45.4
L387A	44.8
S359T/S360T	48.0

HPTLC screening

Reaction samples were analyzed by high performance thin layer chromatography (HPTLC; *CAMAG Automatic TLC Sampler 4*) on Merck silica gel plates 60 F₂₅₄. For each time point measurement two identical 10 µL-samples were sprayed and developed with different solvent mixtures to discriminate partially overlapping bands and quantify the amount of product (α 2,3-SiaLac) and hydrolysis side product (Neu5Ac). The basic solvent mix consisted of *n*-propanol : H₂O : NH₃(20%) = 7 : 3 : 1, and acidic solvent mix of *n*-butanol : acetone : AcOH : H₂O = 35 : 35 : 7 : 23. After anisaldehyde staining a quantitative analysis was performed (*CAMAG TLC Scanner 4*) against standards for SiaLac and Neu5Ac as positive controls, resulting in absolute concentrations of the substrates and products.




Figure S3. Elution by *n*-propanol:H₂O:NH₃ = 7:3:1 for unbiased integration of Neu5Ac.




Figure S4. Quantitative analysis of sample from Figure S3; inset shows integration of lane 2.




Figure S5. Elution by *n*-butanol:acetone:H₂O:AcOH = 35:35:23:7 for unbiased integration of SiaLac.




Figure S6. Quantitative analysis for lane 1.

Table S8. HPTLC analysis of sialyltransfer vs. hydrolysis using 2,3SiaT_{pph} (NE) and variants (duplicate runs).

1.	NE	C8 (A151D)	C7 (L387A)	G7 (S359T/S360T)	2x (A151D/L387A)			
	product %	hydrolysis %	product %	hydrolysis %	product %	hydrolysis %	product %	hydrolysis %
10,00	9,36	2,99	14,43	0,57	7,18	1,31	2,65	3,91
45,00	16,87	2,40	22,64	2,85	14,79	3,30	6,89	2,25
120,00	27,85	2,33	31,73	2,10	25,57	2,73	19,57	2,38
240,00	50,38	7,97	55,82	5,52	44,94	6,89	39,48	4,75
360,00	70,76	6,18	73,50	3,58	65,20	3,91	68,11	3,66
1320,00	89,29	10,71	93,03	6,97	88,11	8,56	91,41	8,59
								29,68
								14,57

2.	NE	C8 (A151D)	C7 (L387A)	G7 (S359T/S360T)	2x (A151D/L387A)			
	product %	hydrolysis %	product %	hydrolysis %	product %	hydrolysis %	product %	hydrolysis %
10,00	6,96	1,73	8,39	0,60	5,57	1,22	2,90	1,81
45,00	19,46	1,85	22,39	1,02	16,67	1,33	12,48	1,67
120,00	30,62	2,01	34,26	1,09	33,57	1,94	30,65	3,47
240,00	46,10	4,55	47,12	2,14	45,90	2,10	47,47	2,25
360,00	51,86	5,30	55,78	2,05	57,83	1,70	65,08	3,06
1320,00	89,08	10,92	93,95	6,05	91,27	6,49	93,71	9,76
								39,73
								11,26

Average	NE	C8 (A151D)	C7 (L387A)	G7 (S359T/S360T)	2x (A151D/L387A)			
	product %	hydrolysis %	product %	hydrolysis %	product %	hydrolysis %	product %	hydrolysis %
10,00	8,16	2,36	11,41	0,58	6,38	1,27	2,77	2,86
45,00	18,17	2,13	22,51	1,94	15,73	2,32	9,68	1,96
120,00	29,24	2,17	33,00	1,60	29,57	2,34	25,11	2,92
240,00	48,24	6,26	51,47	3,83	45,42	4,49	43,48	3,50
360,00	61,31	5,74	64,64	2,82	61,51	2,80	66,59	3,36
1320,00	89,19	10,81	93,49	6,51	89,69	7,53	92,56	9,18
								34,70
								12,92

error	wt	C8 (A151D)	C7 (L387A)	G7 (S359T/S360T)	2x (A151D/L387A)			
	product %	hydrolysis %	product %	hydrolysis %	product %	hydrolysis %	product %	hydrolysis %
10	1,20	0,63	3,02	0,02	0,81	0,04	0,12	1,05
45	1,29	0,27	0,13	0,91	0,94	0,98	2,80	0,29
120	1,39	0,16	1,26	0,51	4,00	0,40	5,54	0,54
240	2,14	1,71	4,35	1,69	0,48	2,40	4,00	1,25
360	9,45	0,44	8,86	0,76	3,68	1,11	1,52	0,30
1320	0,11	0,11	0,46	0,46	1,58	1,04	1,15	0,58
								5,02
								1,65

NMR study of sialidase activity

Table S9. Data for Figure 4: Quantification of sialidase activity of 2,3SiaT_{pph} and variants by ¹H NMR (500 MHz) *in situ* monitoring of Neu5Ac liberation from α2,3-SiaLac.^[a]

	Native SiaT	A151D	L387A	S359T/S360T
18h	10.40	0.48	0.29	0.27
20h	11.72	0.50	0.32	0.38
22h	13.13	0.55	0.33	0.40
24h	13.23	0.60	0.34	0.46
42h	19.16	0.83	0.48	0.71
48h	20.45	0.92	0.65	0.66

[a] Conditions: 14.9 mM α2,3-SiaLac, 520 μL Tris-buffered (20 mM) D₂O, 10 μg enzyme, 30°C, 48h.

Control for CMP independent sialidase activity





Figure S7. Control experiment for sialidase activity in the presence of alkaline phosphatase (Php) by *in situ*-NMR spectroscopy demonstrating hydrolytic Neu5Ac release from GM3 in the absence of CMP. Conditions: 14.9 mM α2,3-SiaLac, 520 μL Tris-buffered (20 mM) D₂O, 10 μg enzyme, 30°C, 48h. Control: + 20 U alkaline phosphatase. NE = native enzyme.

General protocol for synthesis of fluorescence-labeled *neo*-sialoconjugates using native 2,3SiaT_{pph}

CTP (25.2 mg; 2 eq.) and the corresponding sialic acid (Neu5Ac **1** or analogs **2-6**; 1.5 eq.) were dissolved in 5 mL Tris-HCl buffer (50 mM, pH 8.6) containing 0.5 M NaCl, 0.1% Triton×100 and 20 mM MgCl₂. Reaction was started by addition of 5 mg CSS from *N. meningitis* and inorganic pyrophosphatase (2 U). When CMP activation was completed, lactoside **7** (20 mg; 1 eq.), 0.5 mg recombinant 2,3SiaT_{pph} and alkaline phosphatase (160 U) were added and the pH was readjusted. When the reaction was finished, 30 mL cold methanol (−20 °C) was added to stop the reaction. Protein precipitate was removed by filtration. The filtrate was passed over a plug of reverse-phase C₁₈-silica and the product eluted with 20-50% gradient of aqueous methanol. The solution was concentrated to 10 mL and separated by Biogel P-2 (Bio-Rad, Germany) column chromatography (3 × 100 cm) using 5 mM Tris buffer (pH 8.5). The product fractions were collected and lyophilized for characterization.





α2,3-Neu5Ac-Lac-Acr (8) from 13.1 mg Neu5Ac; yield 23.7 mg pale yellow solid (84%).

¹H NMR (500 MHz, D₂O) δ = 7.87 (d, *J* = 8.1 Hz, 2H, 17-, 21-H), 7.82 (s, 1H, 7-H), 7.37 (t, *J* = 7.1 Hz, 2H, 15-, 23-H), 6.98 – 6.90 (m, 4H, 24-, 14-, 16-, 22-H), 5.79 – 5.56 (m, 2H, 2-, 3-H), 4.54 (s, 2H, 5-H), 4.52 (d, *J* = 8.1 Hz, 1H, 1"-H), 4.35 (d, *J* = 8.0 Hz, 1H, 1'-H), 4.32 – 4.22 (m, 3H, 1a-, 8-H), 4.22 – 4.16 (m, 1H, 1b-H), 4.16 – 4.11 (m, 1H), 4.04 – 3.98 (m, 3H, 3"-, 2-, 4-H), 3.96 – 3.85 (m, 5H, 4"-, 6"-, 9""a-, 6'a-, 5""-H), 3.82 – 3.50 (m, 13H, 6'b-, 6"-, 8"-, 7"-, 9""b-, 3'-, 4"-, 4'-, 2"-, 8-, 5"-H), 3.42 (ddd, *J* = 9.6, 4.7, 2.5 Hz, 1H, 5'-H), 3.29 (dd, *J* = 9.1, 8.1 Hz, 1H, 2'-H), 2.81 (dd, *J* = 12.4, 4.6 Hz, 1H, 3""a-H), 2.07 (s, 3H, 11""-H), 1.84 (t, *J* = 12.1 Hz, 1H, 3""b-H), 1.67 – 1.57 (m, 2H, 9-H), 1.19 – 0.99 (m, 4H, 11-, 10-H).

¹³C NMR (126 MHz, D₂O) δ = 178.00 (C-19), 175.03 (C-10""), 173.90 (C-1""), 143.81 (C-6), 140.53 (C-13, -25), 134.49 (C-15, -23), 129.25 (C-2), 129.00 (C-3), 126.19 (C-17, -21), 124.59 (C-7), 121.58 (C-16, -22), 120.34 (C-18, -20), 114.92 (C-14, -24), 102.75 (C-1"), 101.10 (C-1'), 99.83 (C-2""), 78.34 (C-4'), 75.55 (C-3"), 75.15 (C-5"), 74.76 (C-5'), 74.41 (C-3'), 72.93 (C-2'), 72.71 (C-8""), 71.77 (C-6""), 69.37 (C-2"), 68.34 (C-7""), 68.14 (C-4""), 67.50 (C-4"), 65.13 (C-4), 64.70 (C-1), 62.64 (C-9""), 62.32 (C-5), 61.03 (C-6'), 51.77 (C-6"), 50.02 (C-8), 45.20 (C-12), 39.70 (C-3""), 28.80 (C-9), 25.73 (C-11), 22.61 (C-10), 22.08 (C-11").

HRMS ESI⁺: *m/z* calcd for [C₄₈H₆₆N₅O₂₁]⁺ = 1048.4245; found 1048.4255.







α2,3-Neu5Gc-Lac-Acr (9) from 13.7 mg Neu5Gc; yield 22.1 mg pale yellow solid (77%).

¹H NMR (500 MHz, D₂O) δ = 7.95 (dd, *J* = 8.1, 1.6 Hz, 2H, 17-, 21-H), 7.86 (s, 1H, 7-H), 7.46 (ddd, *J* = 8.7, 6.9, 1.6 Hz, 2H, 15-, 23-H), 7.10 – 6.96 (m, 4H, 24-, 14-, 16-, 22-H), 5.84 – 5.63 (m, 2H, 2-, 3-H), 4.59 (s, 2H, 5-H), 4.55 (d, *J* = 7.8 Hz, 1H, 1"-H), 4.38 (d, *J* = 8.0 Hz, 1H, 1'-H), 4.34 – 4.29 (m, 3H, 1a-, 8-H), 4.25 – 4.15 (m, 4H, 11"-, 1b-, 3"-H), 4.07 – 3.79 (m, 12H, 6'a-, 5"-, 6'b-, 6"-, 8""-, 6""-, 9""b-, 3'-, 4""-, 4"-, 2"-H), 3.79 – 3.57 (m, 9H, 6'b-, 3"-, 9""b-, 12-, 7""-, 3'-, 4'-, 2"-H), 3.45 (ddd, *J* = 9.7, 4.9, 2.5 Hz, 1H, 5'-H), 3.32 (dd, *J* = 9.2, 8.0 Hz, 1H, 2'-H), 2.90 – 2.82 (m, 1H, 3""a-H), 1.88 (td, *J* = 12.1, 3.5 Hz, 1H, 3""b-H), 1.69 (p, *J* = 7.0 Hz, 2H, 9-H), 1.20 – 1.06 (m, 4H, 11-, 10-H).

¹³C NMR (126 MHz, D₂O) δ = 178.17 (C-19), 175.82 (C-10""), 173.95 (C-1""), 143.87 (C-6), 140.65 (C-13, -25), 134.60 (C-15, -23), 129.27 (C-2), 129.05 (C-3), 126.25 (C-17, -21), 124.66 (C-7), 121.68 (C-16, -22), 120.41 (C-18, -20), 115.05 (C-14, -24), 102.80 (C-1"), 101.13 (C-1'), 99.89 (C-2""), 78.42 (C-4'), 75.60 (C-3"), 75.19 (C-8""), 74.80 (C-5'), 74.45 (C-5"), 72.74 (C-3'), 72.70 (C-2'), 71.86 (C-6""), 69.41 (C-2"), 68.13 (C-7""), 67.53 (C-4""), 65.10 (C-4), 64.73 (C-1), 62.65 (C-9""), 62.35 (C-5), 61.08 (C-6", -11""), 60.10 (C-6'), 51.51 (C-5""), 50.09 (C-8), 45.24 (C-12), 39.81 (C-3""), 28.81 (C-9), 25.79 (C-11), 22.66 (C-10).

HRMS ESI⁺: *m/z* calcd for [C₄₈H₆₆N₅O₂₂]⁺ = 1064.4194; found 1064.4206.







α2,3-KDN-Lac-Acr (10) from 11.5 mg KDN; yield 23.4 mg pale yellow solid (86%).

^1H NMR (500 MHz, D_2O) δ = 7.92 (dd, J = 8.0, 1.7 Hz, 2H, 17-, 21H), 7.84 (s, 1H, 7-H), 7.45 – 7.35 (m, 2H, 15-, 23-H), 7.05 – 6.92 (m, 4H, 14-, 24-, 16-, 22-H), 5.81 – 5.62 (m, 2H, 2-, 3-H), 4.56 (s, 2H, 5-H), 4.55 – 4.52 (m, 1H, 1"-H), 4.38 (d, J = 8.0 Hz, 1H, 1'-H), 4.35 – 4.25 (m, 3H, 1a-, 8-H), 4.22 (dd, J = 12.6, 6.8 Hz, 1H, 1b-H), 4.17 – 4.13 (m, 1H, H3''), 4.07 – 4.01 (m, 3H), 4.00 – 3.90 (m, 4H), 3.85 – 3.73 (m, 5H), 3.72 – 3.54 (m, 8H, 12-, 9'''b-, 4'''-, 8'''-, 3'-, 2'-, 4'-H), 3.45 (ddd, J = 9.7, 4.8, 2.5 Hz, 1H, 5'-H), 3.42 (s, 1H), 3.32 (dd, J = 9.2, 8.0 Hz, 1H, 2'-H), 2.79 (dd, J = 12.5, 4.6 Hz, 1H, 3'''a-H), 1.83 (td, J = 12.0, 2.8 Hz, 1H, 3'''b-H), 1.69 – 1.60 (m, 2H, 9-H), 1.24 – 1.02 (m, 4H, 11-, 10-H).

^{13}C NMR (126 MHz, D_2O) δ = 178.05 (C-19), 174.11 (C-1''), 143.88 (C-6), 140.61 (C-13, -25), 134.53 (C-15, -23), 129.29 (C-2), 129.04 (C-3), 126.25 (C-17, -21), 124.62 (C-7), 121.61 (C-16, -22), 120.43 (C-18, -20), 114.97 (C-14, -24), 102.82 (C-1''), 101.17 (C-1'), 99.87 (C-2''), 78.43 (C-4'), 75.56 (C-3''), 75.20 (C-8''), 74.79 (C-5'), 74.46 (C-5''), 73.99 (C-3'), 72.75 (C-2'), 72.14 (C-6''), 70.34 (C-2''), 69.85 (C-9''), 62.37 (C-5), 61.06 (C-6''), 60.11 (C-10), 50.05 (C-8), 45.24 (C-12), 39.39 (C-3''), 28.83 (C-9), 25.78 (C-11), 22.66 (C-10).

HRMS ESI $^+$: m/z calcd for $[\text{C}_{46}\text{H}_{63}\text{N}_4\text{O}_{21}]^+$ = 1007.3979; found 1007.3989.







α2,3-*epi*-KDN-Lac-Acr (11) from 11.5 mg *epi*-KDN; yield 15.8 mg pale yellow solid (58%).

¹H NMR (500 MHz, D₂O) δ = 7.94 (dd, *J* = 8.0, 1.7 Hz, 2H, 17-, 21-H), 7.85 (s, 1H, 7-H), 7.51 – 7.38 (m, 2H, 15-, 23-H), 7.08 – 6.95 (m, 4H, 24-, 14-, 16-, 22-H), 5.80 – 5.62 (m, 2H, 2-, 3-H), 4.57 (s, 2H, 5-H), 4.55 (d, *J* = 7.7 Hz, 1H, 1"-H), 4.38 (d, *J* = 8.0 Hz, 1H, 1'-H), 4.32 – 4.28 (m, 3H, 1a-, 8-H), 4.25 – 3.97 (m, 9H, H-3''), 3.96 – 3.56 (m, 16H), 3.46 (ddd, *J* = 9.7, 4.9, 2.5 Hz, 1H, 5'-H), 3.32 (dd, *J* = 9.3, 8.0 Hz, 1H, 2'-H), 2.56 (dd, *J* = 12.2, 4.6 Hz, 1H, 3'''a-H), 2.04 (t, *J* = 12.4 Hz, 1H, 3'''b-H), 1.67 (p, *J* = 7.1 Hz, 2H, 9-H), 1.24 – 1.05 (m, 4H, 11-, 10-H).

¹³C NMR (126 MHz, D₂O) δ = 178.12 (C-19), 174.22 (C-1''), 143.88 (C-6), 140.64 (C-13, -25), 134.56 (C.15, -23), 129.27 (C-2), 129.03 (C-3), 126.26 (C-17, -21), 124.63 (C-7), 121.65 (C-16, -22), 120.42 (C-18, -20), 115.02 (C-14, -24), 102.71 (C-1''), 101.16 (C-1'), 100.13 (C-2''), 78.33 (C-4'), 75.54 (C-3''), 75.22 (C-8''), 74.80 (C-5'), 74.44 (C-5''), 73.02 (C-3'), 72.75 (C-2'), 71.88 (C-6''), 71.68 (C-7'), 71.44 (C-10'), 70.94 (C-11'), 69.55 (C-12'), 69.14 (C-13'), 67.47 (C-14'), 65.13 (C-15'), 64.75 (C-16'), 62.35 (C-17'), 62.18 (C-18'), 61.08 (C-19'), 60.12 (C-20'), 50.07 (C-21'), 45.23 (C-22'), 34.09 (C-23'), 28.82 (C-24'), 25.79 (C-25'), 22.66 (C-26').

HRMS ESI⁺: *m/z* calcd for [C₄₆H₆₃N₄O₂₁]⁺ = 1007.3979; found 1007.3983.







α 2,3-Neu5Ac9N₃-Lac-Acr (12) from 14.1 mg Neu5Ac9N₃; yield 23.7 mg pale yellow solid (82%).

¹H NMR (500 MHz, D₂O) δ = 7.96 – 7.90 (m, 2H, 17-, 21-H), 7.84 (s, 1H, 7-H), 7.44 – 7.36 (m, 2H, 15-, 23-H), 7.04 – 6.93 (m, 4H, 24-, 14-, 16-, 22-H), 5.81 – 5.63 (m, 2H, 2-, 3-H), 4.56 (s, 2H, 5-H), 4.51 (d, J = 7.9 Hz, 1H, 1''-H), 4.38 (d, J = 7.9 Hz, 1H, 1'-H), 4.35 – 4.19 (m, 4H, 1'-, 1a-, 8-H), 4.17 – 4.11 (m, 1H, H-3''), 4.10 – 3.99 (m, 5H, 6'''-, 4-, 6'a-, 8'''-H), 3.95 – 3.87 (m, 2H, 5'''-, 6'a-H), 3.84 – 3.56 (m, 15H), 3.50 (dd, J = 13.2, 6.6 Hz, 1H, 9'''b-H), 3.46 – 3.42 (m, 1H, 5'-H), 3.31 (dd, J = 9.1, 8.0 Hz, 1H, 2'-H), 2.84 (dd, J = 12.4, 4.6 Hz, 1H, 3'''a-H), 2.10 (s, 3H, 11'''-H), 1.85 (t, J = 12.1 Hz, 1H, 3'''b-H), 1.65 (t, J = 7.4 Hz, 2H, 9-H), 1.23 – 1.05 (m, 4H, 11-, 10-H).

¹³C NMR (126 MHz, D₂O) δ = 178.03 (C-19), 175.05 (C-10'''), 173.80 (C-1''), 143.88 (C-6), 140.62 (C-13, -25), 134.51 (C-15, -23), 129.30 (C-2), 129.04 (C-3), 126.29 (C-17, -21), 124.59 (C-7), 121.59 (C-16, -22), 120.47 (C-18, -20), 114.95 (C-14, -24), 102.80 (C-1''), 101.12 (C-1'), 99.83 (C-2'''), 78.36 (C-4'), 75.75 (C-3''), 75.21 (C-8'''), 74.82 (C-5'), 74.45 (C-2''), 70.44 (C-2'), 69.36 (C-6'''), 68.94 (C-2''), 68.30 (C-1''), 67.46 (C-14), 65.17 (C-13), 64.69 (C-12), 62.38 (C-11), 61.04 (C-10), 60.12 (C-9), 59.36 (C-8), 58.29 (C-7), 57.21 (C-6), 56.29 (C-5), 55.17 (C-4), 54.14 (C-3), 53.92 (C-2), 52.48 (C-1), 45.24 (C-10), 42.46 (C-9), 39.92 (C-8), 38.86 (C-7), 38.63 (C-6), 38.44 (C-5), 38.21 (C-4), 38.01 (C-3), 37.79 (C-2), 37.58 (C-1), 36.68 (C-10), 36.66 (C-9), 36.65 (C-8), 36.65 (C-7), 36.64 (C-6), 36.63 (C-5), 36.62 (C-4), 36.60 (C-3), 36.60 (C-2), 35.99 (C-11), 35.58 (C-10), 35.52 (C-9), 35.51 (C-8), 35.50 (C-7), 34.42 (C-6), 33.33 (C-5), 33.31 (C-4), 33.31 (C-3), 32.10 (C-2), 1.85 (C-1), 1.65 (C-10), 1.10 (C-9), 1.09 (C-8).

HRMS ESI⁺: *m/z* calcd for [C₄₈H₆₅N₈O₂₀]⁺ = 1073.4310; found 1073.4321.






α 2,3-Neu5NPhAc-Lac-Acr (13) from 16.1 mg Neu5NPhAc; yield 14.5 mg pale yellow solid (48%).

¹H NMR (500 MHz, D₂O) δ = 7.98 (dd, *J* = 7.9, 1.7 Hz, 2H, 17-, 21-H), 7.84 (s, 1H, 7-H), 7.45 (dt, *J* = 10.7, 8.3 Hz, 2H, 15-, 23-H), 7.42 – 7.28 (m, 5H, Har), 7.11 – 6.99 (m, 4H, 24-, 14-, 16-, 22-H), 5.82 – 5.60 (m, 2H, 2-, 3-H), 4.57 (s, 2H, 5-H), 4.53 (d, *J* = 7.8 Hz, 1H, 1"-H), 4.37 (d, *J* = 7.9 Hz, 1H, 1'-H), 4.33 – 4.25 (m, 3H, 1a-, 8-H), 4.22 (dd, *J* = 12.6, 6.5 Hz, 1H, 1b-H), 4.18 – 4.12 (m, 1H, 3"-H), 4.07 – 3.99 (m, 3H), 3.94 – 3.55 (m, 20H), 3.45 – 3.40 (m, 2H, 7"-, 5'-H), 3.31 (dd, *J* = 9.2, 8.0 Hz, 1H, 2'-H), 2.88 – 2.81 (m, 1H, 3'"a-H), 1.85 (td, *J* = 11.8, 3.7 Hz, 1H, 3'"b-H), 1.68 (p, *J* = 6.3 Hz, 2H, 9-H), 1.29 – 1.06 (m, 4H, 11-, 10-H).


¹³C NMR (126 MHz, D₂O) δ = 178.21 (C-19), 175.60 (C-10''), 173.88 (C-1'''), 143.85 (C-6), 140.71 (C-13, -25), 135.04 (C-15, -23), 134.61 (C-12''), 129.34 (C-14'''), 129.09 (C-2), 129.01 (C-3), 128.91 (C-13''', -15''), 127.30 (C-12'', -16''), 126.29 (C-17, -21), 124.64 (C-7), 121.69 (C-16, -22), 120.48 (C-18, -20), 115.08 (C-14, -24), 102.82 (C-1''), 100.99 (C-1'), 99.81 (C-2''), 78.35 (C-4'), 75.62 (C-3''), 75.21 (C-8''), 74.79 (C-5'), 74.44 (C-5''), 72.99 (C-3'), 72.74 (C-2'), 71.85 (C-6''), 69.37 (C-2''), 68.32 (C-7''), 67.53 (C-4'''), 65.08 (C-4), 64.60 (C-1), 62.80 (C-9''), 62.35 (C-5), 61.05 (C-6''), 60.03 (C-6'), 59.84 (C-12), 51.92 (C-5'''), 50.08 (C-8), 42.60 (C-11''), 39.95 (C-3'''), 28.84 (C-9), 25.82 (C-11), 22.69 (C-10).

HRMS ESI⁺: *m/z* calcd for [C₅₄H₇₀N₅O₂₁]⁺ = 1124.4558; found 1124.4572.




Synthesis of α 2,3-SiaLac (GM3 trisaccharide, 14) by native 2,3SiaT_{pph} and engineered 2,3SiaT_{pph} (A151D)


Neu5Ac (50 mg, 1 eq.) and CTP (117.2 mg, 1.5 eq.) were dissolved in 12 mL Tris-HCl buffer (50 mM, pH 8.5) containing 20 mM MgCl₂. After addition of inorganic pyrophosphatase (10 U) the reaction mixture was re-adjusted to pH 8.5 and started by adding 10 mg CSS, while keeping the pH constant by autotitration. After 2 h at 37 °C TLC analysis indicated the disappearance of Neu5Ac, then lactose (174.7 mg, 3 eq.), native 2,3SiaT_{pph} or 2,3SiaT_{pph} (A151D) (0.5 mg) and alkaline phosphatase (319 U) were added and incubation continued overnight. When conversion was completed, an equivalent volume of cold methanol (−20 °C) was added to stop the reaction. Protein precipitate was removed by centrifugation, methanol evaporated under reduced pressure, and the remaining solution directly purified over Biogel P-2 (Bio-Rad, Germany; 3 × 100 cm column). The product fractions were collected and lyophilized for NMR analysis. Product **14** was obtained as colorless solid (88.2 mg, 86% for native 2,3SiaT_{pph} / 93.4 mg, 91% for 2,3SiaT_{pph} (A151D)).



¹H NMR (500 MHz, D₂O) δ = 5.29 (d, *J* = 3.7 Hz, 0.4H, 1α-H), 4.73 (d, *J* = 8.1 Hz, 0.6H, 1β-H), 4.59 (d, *J* = 7.9 Hz, 1H, 1'β-H), 4.18 (ddd, *J* = 9.9, 1.9, 2.9 Hz, 1H, 3'-H), 4.06-3.99 (m, 1.8H, 4'-, 6α-H), 3.98-3.85 (m, 5H, 6'', 5'', 5-, 6'-H), 3.82-3.74 (m, 4H, 6β-, 5'-, 4-H) 3.74-3.68 (m, 4H, 4'', 3-, 9''-H), 3.68-3.65 (m, 2H, 7'', 8''-H), 3.64-3.61 (m, 1H, 2'-H), 3.35 (dd, *J* = 8.5 Hz, 1H, 2-H), 2.82 (dd, *J* = 12.4, 4.6 Hz, 1H, 3''-H), 2.10 (s, 3H, 11''-H), 1.86 (dd, *J* = 12.4, 12.1 Hz, 1H, 3''-H).


¹³C NMR (126 MHz, D₂O) δ = 175.08 (C-1''), 173.90 (C-10''), 102.71 (C-1'), 99.89 (C-2''), 95.83 (C-1β), 91.88 (C-1α), 78.46(C-4α), 78.32 (C-4β), 75.56 (C-3'), 75.21 (C-5'), 74.85(C-2α), 74.39 (C-2β), 73.89 (C-4''), 72.94 (C-3), 71.83 (C-6''), 71.44 (C-5α), 71.23 (C-5β), 69.43 (C-2'), 68.37(C-8''), 68.19 (C-7''), 67.57 (C-4'), 62.68 (C-9''), 61.08 (C-6'), 60.18 (C-6α), 59.48 (C-6β), 51.77(C-5''), 39.70 (C-3''), 22.14 (C-11'').






Exemplary kinetic profiles for native enzyme (NE) and best variants


Enzymatic assays were performed as described. Experimental data were fitted to the Michaelis–Menten equation by nonlinear regression using Origin software (v9.1G, OriginLab) to derive apparent kinetic parameters.




Quality control of second-generation combination library by sequencing

Because positions 148, 150 and 151 are too close to each other to assure sufficient homogenous primer binding, positions 148-151 were first deleted with adjacent primers. Into the resulting template the desired mutated sequence was introduced by using appropriate overhang primers. The PCR product resulting from the first rounds contained the newly introduced sequence, which was used as template for further amplification cycles. The resulting plasmid mixes were sequenced to assure successful mutagenesis. For ambiguous sequence data single clones were sequenced to assure a positive result. All combinations were found:






336/337 muA single clones




muC 43DA00


148 TCG/TTG (Ser/Leu), 150 AGC/ACC (Ser/Thr),
151 GCC (Ala)

**muC 35GH09**


148Ser, 150Ser, 151Ala

**muC 35GH10**


148 Ser, 150 Ser, 151 Ala

**muC 35GH11**


148 Leu, 150 Ser, 151 Ala

**muD 43DA01**

148 GAT (Asp), 150 AGC/ACC (Ser/Thr), 151 GCC (Ala)


**muD single clone 35GH12**

148 GAT (Asp), 150 ACC (Thr), 151 GCC (Ala)




muD single clone 35GH13

148 GAT (Asp), 150 AGC (Ser), 151 GCC (Ala)




muD single clone 35GH14

148 Asp, 150 Thr, 151 Ala



35GH46 muD single clone

148 Asp, 150 Thr, 151 Ala



muD single clone 35GH47

148 GAT (Asp), 150 AGC (Ser), 151 GCC (Ala)

