





Azulene Renaissance

The highly symmetric 2,4,6,8-tetracyanoazulene is obtained by stepwise substitution of the seven-membered ring of 2-cyanoazulene by nitrile groups.

It joins the series of strong acceptors (TCNE, TCNQ, and DCNQI) of electrically conductive charge-transfer complexes and radical anion salts.




The synthesis and properties of this novel acceptor are reported by K. Hafner et al. on the following pages.

2,4,6,8-Tetracyanoazulene: A New Building Block for "Organic Metals"***

Stefan Schmitt, Martin Baumgarten, Joachim Simon, and Klaus Hafner*

Organic charge-transfer (CT) complexes are being increasingly investigated, both theoretically and experimentally,^[1] as their molecular structure allows an easy preparation of single-crystal, well-characterizable compounds with high electric conductivity. The formation of such CT complexes is based on the interaction of an energetically high-lying HOMO (of the donor) with a low-energy LUMO (of the acceptor). Many one-dimensional compounds derived from the acceptors 7,7,8-tetracyano-*p*-quinodimethane (TCNQ, **1**)^[2] and *N,N'*-dicyano-*p*-quinodimine (DCNQI, **2**)^[3] as well as the donor tetrathiafulvalene (TTF, **3**; Scheme 1)^[4] have already been investigated; superconductivity was proven even for radical ion salts.^[5]



Scheme 1. Electron acceptors and donors.

The transfer of free electrons along separated donor or acceptor stacks can formally be understood as "migration of aromaticity".^[6] Accordingly, resonance structures with a cyclic electron sextet can be formulated for the radical ions of **1–3**.





Figure 1. Schematic representation of the radical anion **4A** and the radical cation **4B** as well as the LUMO coefficients of azulene (**4**).

As numerous experimental studies and calculations^[7] of the redox characteristics of azulene (**4**)—a bicyclic, non-alternating hydrocarbon—and its derivatives have shown, its radical anion **4A** (cyclopentadienide ion) and cation **4B** (tropylium cation) can be described with an aromatic partial structure (Figure 1). Whereas the donor properties of azulene have been thoroughly inves-

tigated,^[8] none of its substituted derivatives has been employed so far as an acceptor for the preparation of CT complexes. According to HMO theory, the incorporation of acceptor substituents at the azulene positions with large LUMO coefficients (2-, 4-, 6-, and 8-positions; Figure 1) should lead to a decrease in the energy of the LUMO and consequently to an increased electron affinity.^[9]

We report here on the synthesis of 2,4,6,8-tetracyanoazulene (TCNA, **10**) and the di- and tricyanoazulenes **6–9** (Scheme 2) as well as their chemical and physical properties. Our special interest was focused on the electron affinity of these new π acceptors and their tendency to form CT complexes.



Scheme 2. Synthesis of azulenenitriles **6–10**. a) 1. KCN/DMF; 2. 2 N AcOH; 3. *p*-chloranil.

In contrast to azulene (**4**), which can be attacked only by strong nucleophiles such as alkali organometallic compounds at the 4- or 8- and 6-positions,^[10] azulenes with acceptor substituents on the five-membered ring react with cyanide ions in dimethylformamide (DMF) already at room temperature to form σ adducts.^[11] After hydrolysis with 2 N AcOH and dehydrogenation with *p*-chloranil, these adducts can be converted into 4-, 8-, and 6-cyanoazulenes as blue or green needles. 2,4,6,8-Tetracyanoazulene (**10**) was thus obtained from 2-cyanoazulene (**5**)^[12] by repeated nucleophilic addition and dehydrogenation.^[13] The ratios of regioisomers are in agreement with findings by McDonald et al.,^[14] according to which the reaction of azulenes with nucleophiles affords mixtures of regioisomers providing an attack at one of these positions is not favored because of steric or electronic reasons. Owing to the differences in symmetry, differentiation between

[*] Prof. Dr. K. Hafner, Dipl.-Ing. S. Schmitt
 Institut für Organische Chemie der Technischen Universität
 Petersenstrasse 22, D-64287 Darmstadt (Germany)
 Fax: (+49) 6151-163574
 E-mail: khafner@hrz.tu.darmstadt.de
 Priv.-Doz. Dr. M. Baumgarten
 Max-Planck-Institut für Polymerforschung, Mainz (Germany)
 Dr. J. Simon
 Institut für Anorganische Chemie der
 Technischen Universität Darmstadt

[**] This work was supported by the Fonds der Chemischen Industrie, the Dr.-Otto-Röhm-Gedächtnisstiftung, and Degussa AG.

compounds **6** and **7** as well as between **8** and **9** is possible by ^1H NMR spectroscopy (Scheme 2).

Characteristic for the substitution at an electron-deficient position of the seven-membered ring by a nitrile group is a bathochromic shift of the long-wave absorption of azulenes **6–10**. This suggests a decrease in the energy of the LUMO. Additional information on the energy of the frontier orbitals as well as the electron affinity of compounds **6**, **8**, and **10** was obtained by cyclic voltammetry studies. These indicate that the formation of the radical anions and the paratropic dianions is completely reversible (Figure 2). A comparison of the half-wave potentials for the first reduction of **6**, **8**, and **10** shows




Figure 2. Cyclic voltammogram of **10** ($n\text{Bu}_4\text{NPF}_6$, 10^{-4}M , CH_2Cl_2 , 25°C , 100mVs^{-1} , working electrode: Pt disk, counter electrode: Pt plate (1cm^2), reference electrode: saturated calomel electrode (SCE)).

that the half-wave potential increases with rising substitution at the electron-deficient positions of the seven-membered ring (Table 1). These data are in agreement with a resonance stabilization of the negative charge and the unpaired electron by the cyano groups.

Table 1. First and second half-wave potentials of azulenenitriles **6**, **8**, and **10**.

Compd	E_1 [mV]	E_2 [mV]
6	-765	-1631
8	-320	-1240
10	80	-795

First indications for the ability of **10** to form colored CT complexes were provided by UV/Vis studies on solutions of **10** and N,N,N',N' -tetramethyl-*p*-phenylenediamine (TMPD) in dichloromethane or acetonitrile (Figure 3). Whereas in dichloromethane two unstructured, concentration-dependent CT bands at 513 and 1200 nm appear in addition to the characteristic absorptions of **10** between 600 and 950 nm, in acetonitrile absorption bands are observed that indicate the presence of the Wurster radical $\text{TMPD}^{+\cdot}$ and thus indicate a complete charge transfer.

The concentration of CT complexes in solution and the degree of charge transfer are determined with the complex-formation constant and with the difference in potentials





Figure 3. UV/Vis spectra of **10**/TMPD in CH_2Cl_2 (—) and CH_3CN (---). The extinction E is given in arbitrary units.

between the acceptor and donor. Consequently, a more positive reduction or oxidation potential leads to a stronger saltlike character and therefore to lower solubility in organic solvents.^[15] The reaction of **10** with an excess of potassium iodide in acetonitrile yields the potassium salt of the radical anion **11** as a green powder; it is stable in solution and in the solid state at room temperature as well as in the presence of oxygen, and dissolves in polar solvents with an intense violet color (Scheme 3).



Scheme 3. Synthesis of the radical anion salt **11**.

The UV/Vis spectrum of **11** in acetonitrile shows three absorption maxima at $\lambda = 489, 517$ and 554nm together with a broad band of low extinction with characteristic fine structure between 600 and 1100 nm. Information on the distribution of the spin density at the individual positions was anticipated from the distances of the corresponding multiplets in the ESR spectrum. Numerous lines (about 100) appear in the ESR spectrum (Figure 4 a); however, owing to the small width of




Figure 4. ESR spectra of **11**: a) in CH_3CN ; b) simulated.

the signal, they could not be assigned. The proton hyperfine coupling constants of 0.95 and 0.45 G were clearly determined with the ENDOR technique. The nitrogen coupling constants could be established from these by computer simulation^[16]





Figure 5. Nitrogen and proton hyperfine coupling constants [G] of **11**.

tions in the *g* value of the free electron, which are caused by an increase in the spin-orbit coupling value.

Prompted by these findings, it appeared interesting to investigate whether an aprotic, dipolar solvent could also increase the charge transfer of organic donors. In fact, combining solutions of **10** and **3** in acetonitrile provides the CT complex **12** as a microcrystalline powder; the 1:1 ratio of the components was confirmed by elemental analysis.^[13] Complex **12** has a conductivity of 1–3 Scm⁻¹, as determined by a two-point powder measurement (Scheme 4).^[17]



Scheme 4. Synthesis of a CT complex from **10** and **3**.

In accordance with the changed bond character of **11** and **12** as compared to **10**, a characteristic decrease in the CN wavenumber, which is often used as a measure of charge transfer, is observed in the respective IR spectra.

Experimental Section

Standard procedure for the preparation of azulenenitriles: Under an atmosphere of pure nitrogen, the azulene (1.0 mmol) substituted with an acceptor group at the five-membered ring and potassium cyanide (72 mg, 1.1 mmol) are dissolved in anhydrous DMF (10 mL) and stirred at room temperature until the solution turns colorless (ca. 12 h). After subsequent hydrolysis with 2N acetic acid (1 mL), the mixture is poured into a saturated solution of NaCl (100 mL) and extracted with dichloromethane (3 × 50 mL). The organic layer is dried with magnesium sulfate, filtered, and treated with *p*-chloranil (245 mg, 1.0 mmol). After 16 h at room temperature the solvent is removed in vacuo, and the residue subjected to chromatography on silica gel. Unchanged *p*-chloranil and starting material can be recovered in early fractions, and then the desired azulene products substituted at the 4-, 6-, and/or 8-position are isolated.

6: M.p. 174–176 °C; ¹H NMR (300 MHz, CD₂Cl₂, 25 °C, TMS): δ = 7.46 (t, ³J(H,H) = 9.8 Hz, 1H, 7-H), 7.51 (d, ³J(H,H) = 9.9 Hz, 1H, 5-H), 7.70 (d, ³J(H,H) = 1.6 Hz, 1H, 1-H), 7.75 (t, ³J(H,H) = 9.1 Hz, 1H, 6-H), 7.85 (d, ⁴J(H,H) = 1.2 Hz, 1H, 3-H), 8.47 (d, ³J(H,H) = 9.3 Hz, 1H, 8-H); IR (KBr):

$\tilde{\nu}$ = 2229 cm⁻¹ (CN); UV/Vis (CH₂Cl₂): λ_{max} (lg ε) = 257 (3.94), 295 (4.24), 307 (sh, 4.02), 348 (3.13), 362 (3.19), 367 (3.16), 379 (3.12), 597 (sh, 2.18), 636 (2.29), 689 (2.27), 758 nm (1.93); MS (70 eV): *m/z* (%): 178 (100), 151 (19) [*M* – HCN]⁺.

7: M.p. 225 °C (subl.); ¹H NMR (300 MHz, CD₂Cl₂, 25 °C, TMS): δ = 7.47 (d, ³J(H,H) = 10.3 Hz, 1H, 7-H and 5-H), 7.68 (s, 1H, 1-H and 3-H), 8.38 (d, ³J(H,H) = 10.4 Hz, 1H, 4-H and 8-H); IR (KBr): $\tilde{\nu}$ = 2227 cm⁻¹ (CN); UV/Vis (CH₂Cl₂): λ_{max} (lg ε) = 281 (4.76), 327 (sh, 3.49), 338 (3.68), 351 (3.85), 362 (sh, 3.70), 368 (4.06), 603 (sh, 2.55), 646 (2.70), 702 (2.68), 775 nm (2.32); MS (70 eV): *m/z* (%): 178 (100), 151 (26) [*M* – HCN]⁺.

8: M.p. 264–265 °C; ¹H NMR (300 MHz, CD₂Cl₂, 25 °C, TMS): δ = 7.75 (d, ³J(H,H) = 10.2 Hz, 1H, 5-H and 7-H), 7.93 (t, ³J(H,H) = 9.4 Hz, 1H, 6-H), 8.10 (s, 1-H and 3-H); IR (KBr): $\tilde{\nu}$ = 2227 cm⁻¹ (CN); UV/Vis (CH₂Cl₂): λ_{max} (lg ε) = 269 (4.33), 291 (sh, 4.21), 302 (4.30), 311 (4.32), 324 (sh, 4.00), 360 (3.30), 377 (3.39), 395 (3.17), 639 (sh, 2.30), 684 (2.44), 752 (2.42), 840 nm (2.01); MS (70 eV): *m/z* (%): 203 (100), 176 (15) [*M* – HCN]⁺.

9: M.p. 268 °C (subl.); ¹H NMR (300 MHz, CD₂Cl₂, 25 °C, TMS): δ = 7.67 (s, 1H, 5-H), 7.71 (d, ³J(H,H) = 10.2 Hz, 1H, 7-H), 7.85 (s, 1H, 3-H), 7.97 (s, 1H, 1-H), 8.50 (d, ³J(H,H) = 9.7 Hz, 1H, 8-H); IR (KBr): $\tilde{\nu}$ = 2222 cm⁻¹ (CN); UV/Vis (CH₂Cl₂): λ_{max} (lg ε) = 295 (4.72), 353 (3.54), 371 (3.75), 389 (3.87), 637 (sh, 2.49), 684 (2.63), 752 (2.61), 838 nm (2.21); MS (70 eV): *m/z* (%): 203(100), 176 (21) [*M* – HCN]⁺.

10: M.p. 348 °C (decomp); ¹H NMR (300 MHz, CD₂Cl₂, 25 °C, TMS): δ = 7.94 (s, 2H, 5-H and 7-H), 8.24 (s, 2H, 1-H and 3-H); IR (KBr): $\tilde{\nu}$ = 2235 cm⁻¹ (CN); UV/Vis (CH₂Cl₂): λ_{max} (lg ε) = 277 (4.43), 306 (4.78), 366 (3.60), 385 (3.83), 404 (3.82), 675 (sh, 2.49), 725 (2.63), 802 (2.58), 912 nm (2.19); MS (70 eV): *m/z* (%): 228 (100), 201 (20) [*M* – HCN]⁺.

11: M.p. 281 °C (decomp); IR (KBr): $\tilde{\nu}$ = 2197 cm⁻¹ (CN); UV/Vis (CH₃CN): λ_{max} = 247, 306, 325 (sh), 364, 380, 399, 410 (sh), 462 (sh), 489, 517, 554, 687, 753, 864, 1000 nm.

12: M.p. 256 °C (decomp); IR (KBr): $\tilde{\nu}$ = 2201 cm⁻¹ (CN).

Received: October 17, 1997 [Z11043IE]
German version: *Angew. Chem.* **1998**, *110*, 1129–1133

Keywords: azulene • charge transfer • cyclic voltammetry • donor–acceptor systems • EPR spectroscopy

[1] a) M. R. Bryce, *Introd. Mol. Electron.* **1995**, 168–184; b) C. M. Gossel, C. S. Weston, *Contemp. Org. Synth.* **1994**, *1*, 367–385.
[2] a) T. Yanagimoto, K. Takimiya, T. Otsubo, T. Ojura, *J. Chem. Soc. Commun.* **1993**, 519–520; b) Y. Tsubata, T. Suzuki, T. Miyasahi, Y. Yamashita, *J. Org. Chem.* **1992**, *57*, 6749–6755; c) P. Gruz, N. Martin, F. Miguel, C. Seoane, A. Albert, F. H. Cano, A. Gonzalez, J. M. Pingarron, *ibid.* **1992**, *57*, 6192–6198.
[3] a) A. Aumüller, P. Erk, S. Hünig, J. U. von Schütz, H.-P. Werner, H. C. Wolf, G. Klebe, *Chem. Ber.* **1991**, *124*, 1445–1451; b) S. Hünig, *J. Mater. Chem.* **1995**, *5*, 1469–1479.
[4] a) J. M. Williams, A. J. Sultz, U. Geiser, K. D. Carlson, A. M. Kini, H. H. Wang, W.-K. Kwok, M. H. Wangbo, J. E. Schreiber, *Science* **1991**, *252*, 1501–1508; b) M. R. Bryce, *Chem. Soc. Rev.* **1991**, *20*, 355–390.
[5] a) A. Aumüller, P. Erk, G. Klebe, S. Hünig, J. U. von Schütz, H.-P. Werner, *Angew. Chem.* **1986**, *98*, 759–761; *Angew. Chem. Int. Ed. Engl.* **1986**, *25*, 740–742; b) J. M. Williams, J. R. Ferraro, R. J. Thorn, K. D. Carlson, U. Geiser, H. H. Wang, A. M. Kini, M.-H. Whangbo, *Organic Superconductors, Synthesis, Structure, Properties and Theory*, Prentice-Hall, Englewood Cliffs, NJ, USA, **1992**.
[6] J. H. Perlstein, *Angew. Chem.* **1977**, *89*, 534–549; *Angew. Chem. Int. Ed. Engl.* **1977**, *16*, 519–554.
[7] a) R. Bachmann, C. Burda, F. Gerson, M. Scholz, H.-J. Hansen, *Helv. Chim. Acta.* **1994**, *77*, 1458–1465; b) I. Bernal, P. H. Rieger, G. K. Fraenkel, *J. Chem. Phys.* **1962**, *37*, 1489–1495; c) R. M. Dessau, S. Shih, *ibid.* **1970**, *53*, 3169–3172.
[8] a) J. E. Frey, A. M. Andrews, S. D. Combs, S. P. Edens, J. J. Puckett, R. E. Seagle, L. A. Torrealano, *J. Org. Chem.* **1992**, *57*, 6460–6466; b) K. Hafner, K.-L. Moritz, *Justus Liebigs Ann. Chem.* **1961**, *650*, 92–96; c) K. Hafner, K.-L. Moritz, *Angew. Chem.* **1960**, *72*, 918.

- [9] I. Fleming, *Grenzorbitale und Reaktionen organischer Verbindungen*, VCH, Weinheim, 1990.
- [10] a) K. Hafner, H. Weldes, *Justus Liebigs Ann. Chem.* **1957**, 606, 90–99; b) K. Hafner, C. Bernhard, R. Müller, *ibid.* **1961**, 650, 35–39; c) K.-P. Zeller, *Methoden Org. Chem. (Houben-Weyl) 4th ed. 1952–*, Vol. 5/2c, **1985**, pp. 127–418.
- [11] M. Römer, Dissertation, Technische Hochschule Darmstadt, **1977**.
- [12] a) T. Nozoe, S. Seto, S. Matsumura, *Bull. Chem. Soc. Jpn.* **1962**, 35, 1990–1998; b) M. K. O. Goltz, Dissertation, Technische Hochschule Darmstadt, **1986**.
- [13] Correct elemental analyses were obtained for all compounds described here.
- [14] R. N. McDonald, H. E. Petty, N. C. Wolfe, J. V. Paukstelis, *J. Org. Chem.*, **1974**, 39, 1877–1887.
- [15] R. C. Wheland, *J. Am. Chem. Soc.* **1976**, 98, 3926–3930.
- [16] Simulated spectra were obtained with a computer program from the Bruker company for isotropic doublet radicals ($S=1/2$) based on Zeeman and hyperfine interactions. ENDOR spectra enabled the determination of both proton hyperfine couplings used in the simulation. The nitrogen couplings under consideration of symmetry and the determined spin densities were calculated with these computer simulations. The total width of the ESR spectrum already suggested that only small nitrogen couplings would be present: H. Kurreck, B. Kirste, W. Lubitz, *Electron Nuclear Double Resonance Spectroscopy of Radicals in Solution*, VCH, Deerfield Beach, FL, USA, **1988**.
- [17] We thank Dipl.-Ing. J. Leuning (Max-Planck-Institut für Polymerforschung, Mainz, Germany) for the powder conductivity measurements.

Effect of Chain Length on the Adsorption Behavior of *n*-Alkanes in Ferrierite

Willy J. M. van Well, Xavier Cottin, Jan W. de Haan, Rutger A. van Santen, and Berend Smit*

Zeolites are microporous crystalline materials that are used as catalysts in petrochemical applications because of their acidity and stability.^[1] At present there are more than a hundred types of zeolites, each with characteristic pore sizes and pore topologies. One important aim of zeolite science is to use these differences to tune the selectivity of a given reaction.^[2] To arrive at a rational design, it is essential to have an understanding on the molecular level of the behavior of the sorbates in the pores of these zeolites.^[3] Currently such information is mainly obtained from computer simulations. For example, recent Monte Carlo calculations have allowed many predictions to be made about the behavior of *n*-alkanes in zeolites,^[4–8] in particular regarding the preferential adsorption sites of *n*-alkanes in zeolites. To the best of our knowledge, none of these predictions have been confirmed

[*] Prof. Dr. ir. B. Smit
Department of Chemical Engineering
University of Amsterdam
Nieuwe Achtergracht 166, NL-1018 WV Amsterdam
(The Netherlands)
Fax: (+ 31)20-525-5604
E-mail: smit@chemeng.chem.uva.nl
Ir. W. J. M. van Well, Dr. X. Cottin, Dr. ir. J. W. de Haan,
Prof. Dr. R. A. van Santen
Schuit Institute of Catalysis
Eindhoven University of Technology
(The Netherlands)

experimentally, since direct information on the siting of *n*-alkanes is difficult to obtain.

Here we present work on the siting of *n*-alkanes in the zeolite ferrierite (FER). FER has ten-ring channels which are interconnected by cages with eight-ring windows. The use of this zeolite as an isomerization catalyst for the production of isobutene, an intermediate in the production of methyl *tert*-butyl ether (MTBE, considered to be an environmentally beneficial fuel), has been studied extensively.^[9,10] The comparison of the performance of FER and other zeolites as catalysts for the isomerization of *n*-butene has been mainly focused on the dimensions of the ten-ring channels.^[9] We now find that the characteristic pore topology is an important factor as well. One of the questions still open pertains to the distribution of the molecules over these ten- and eight-ring channels.

To study this distribution, we investigated the siting of *n*-alkanes in FER by means of Monte Carlo simulations and ¹³C NMR spectroscopy. For the simulations we assumed that the zeolite is rigid, and the *n*-alkanes were modeled with a united atom representation; that is, CH₃ and CH₂ groups are considered as “united atoms”. Fixed bond lengths, bond angles, and torsion potentials were used.^[11] Details on the simulation techniques have already been described.^[5,12] The results of Monte Carlo simulations (Figure 1) indicate a




Figure 1. Fraction A_{8-Ring} of the molecules adsorbed in the eight-ring channels as a function of the loading a . The data were obtained from computer simulations with the configurational-bias Monte Carlo method at 333 K. ○ propane, □ butane, ◇ pentane, △ hexane.

surprising effect of the chain length. *n*-Alkanes with more than five carbon atoms prefer the ten-ring channels for all loadings; these molecules are too large to fit into the eight-ring channels. At low loadings propane and butane are distributed over the ten-ring and eight-ring channels. While propane prefers adsorption in the eight-ring channel, butane adsorbs preferentially in the ten-ring channel. The adsorption of pentane is at low loading limited to the ten-ring channel, and adsorption in the eight-ring channel occurs only at higher loadings. These simulations show that the adsorption behavior of even simple *n*-alkanes is quite complex and depends strongly on the chain length. One of the purposes of the present work was to present experimental evidence that the above prediction is indeed correct.

An obvious experimental technique to investigate the siting of hydrocarbons in zeolites is ¹³C NMR spectroscopy. The chemical shifts of *n*-alkanes depend on the local environment