

## 7 Literaturverzeichnis

Aigner, A. (1995)

“Transport- und Entgiftungswege an der Blut-Hirn-Schranke: Reinigung, Charakterisierung und Lokalisation der Cystein-S-Konjugat-spezifischen N-Acetyl-S-Transferase des Schweins”

Dissertation, TH-Darmstadt.

Alban, A., David, S. O., Bjorkesten, L., Andersson, C., Sloge, E., Lewis, S. und Currie, I. (2003)

“A novel experimental design for comparative two-dimensional gel analysis: Two-dimensional difference gel electrophoresis incorporating a pooled internal standard”

Proteomics **3**, 36-44.

Allt, G. und Lawrenson, J. G. (2001)

“Pericytes: Cell Biology and Pathology”

Cells Tissues Organs **169**, 1-11.

Anderson, L. und Seilhammer, J. (1997)

“A comparison of selected mRNA and protein abundances in human liver”

Electrophoresis **18**, 533-537.

Baciu, P. C. und Durham, J. P. (1990)

“A procedure for the extraction and high resolution two-dimensional gel electrophoresis of total nuclear phosphoproteins from isotonicity purified nuclei”

Electrophoresis **11**, 162-166.

Bayer, U., De Koninck, P. und Schulman, H. (2002)

“Alternative splicing modulates the frequency-dependent response of CaMKII to Ca<sup>2+</sup> oscillations”

EMBO J. **21**, 3590-3597.

Bayer, U., Harbers, K. und Schulman, H. (1998)

“ $\alpha$ -KAP is an anchoring protein for a novel CaM kinase II isoform in skeletal muscle”

EMBO J. **17**, 5598-5605.

Bayer, U., Löhler, J. und Harbers, K. (1996)

“An alternative, nonkinase product of the brain-specifically expressed Ca<sup>2+</sup>/calmodulin-dependent kinase II a isoform gene in skeletal muscle”

Mol. Cell. Biol. **16**, 29-36.

Benvenuti, S., Cramer, R., Quinn, C. C., Bruce, J., Zvelebil, M., Corless, S., Bond, J., Yang, A., Hockfield, S., Burlingame, A. L., Waterfield, M. D. und Jat, P. S. (2002)

“Differential proteome analysis of replicative senescence in rat embryo fibroblasts”

*Proteomics* **1**, 280-292.

Berggren, K., Chernokalskaya, E., Steinberg, T. H., Kemper, C., Lopez, M. F., Diwu, Z., Haugland, R. P. und Patton, W. F. (2000)

“Background-free, high sensitivity staining of proteins in one- and two-dimensional sodium dodecyl sulfate-polyacrylamide gels using a luminescent ruthenium complex”

*Electrophoresis* **21**, 2509-2521.

Bienvenut, W. V., Sanchez, J. C., Karmime, A., Rouge, V., Rose, K., Binz, P. und Hochstrasser, D. F. (1999)

“Toward a clinical molecular scanner for proteome research: parallel protein chemical processing before and during western blot”

*Anal. Chem.* **71**, 4800-4807.

Binz, P. A., Müller, M., Walther, D., Bienvenut, W.V., Gras, R., Hoogland, C., Bouchet, G., Gasteiger, E., Fabbretti, R., Gay, S., Palagi, P., Wilkins, M. R., Rouge, V., Tonella, L., Paesano, S., Rossellat, G., Karmime, A., Bairoch, A., Sanchez, J. C., Appel, R.D. und Hochstrasser, D. F. (1999).

“A molecular scanner to automate proteomic research and to display proteome images”

*Anal. Chem.* **71**, 4981-4988.

Bjellqvist, B., Sanchez, J. C., Pasquali, C., Ravier, F., Paquet, N., Frutiger, S., Hughes, G. J. und Hochstrasser, D. (1993)

“Micropreparative two-dimensional electrophoresis allowing the separation of samples containing milligram amounts of proteins”

*Electrophoresis* **14**, 1375-1380.

Blum, M., Beier, H. und Gross, H. J. (1987)

“Improved silverstaining of plant proteins, RNA and DNA in polyacrylamide gels”

*Electrophoresis* **8**, 93-99.

Bordier, C. (1981)

“Phase separation of integral membrane proteins in Triton X-114 solution”

*J. Biol. Chem.* **256**, 1604-1607.

Borgese, N., D'Arrigo, A., De Silvestris, M. und Pietrini, G. (1993)

“NADH-cytochrome b5 reductase and cytochrome b5 isoforms as models for the study of post-translational targeting to the endoplasmic reticulum”

FEBS Lett. **325**, 70-75.

Bowman, P. D., Betz, A. L., Ar, D., Wolinsky, J., Penney, J. B., Shivers, R. R. und Goldstein, G. W. (1981)

“Primary culture of capillary endothelium from rat brain”

In Vitro **17**, 353-362.

Bradbury, M. W. (1993)

“The blood-brain barrier”

Exp. Physiol. **78**, 453-472.

Bradford, M. M. (1976)

“A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding”

Anal. Biochem. **72**, 248-254.

Brocke, L., Srinivasan, M., und Schulman, H. (1995)

“Developmental and regional Expression of multifunctional Ca<sup>2+</sup>/calmodulin-dependent protein kinase isoforms in rat brain”

J. Neurosci. **15**, 6797-6808.

Butt, A. M., Jones, H. C. und Abbott, N. J. (1990)

“Electrical resistance across the blood-brain barrier in anaesthetized rats: a developmental study”

J. Physiol. **429**, 47-62.

Cattaruzza, M., Dimigen, C., Ehrenreich, H. und Hecker, M. (2000)

“Stretch-induced endothelin B receptor-mediated apoptosis in vascular smooth muscle cells”

FASEB J. **14**, 991-998.

Chen, M. X., McPartlin, A. E. Brown, L., Chen, Y. H., Barker, H.M. und Cohen, P. T. (1994)

“A novel human protein serine/threonine phosphatase, which possesses four tetratricopeptide repeat motifs and localizes to the nucleus”

EMBO J. **13**, 4278-4290.

Chevallet, M., Santoni, V., Poinas, A., Rouquie, D., Fuchs, A., Kieffer, S., Rossignol, M., Lunardi, J., Garin, J. und Rabilloud, T. (1998)

“New zwitterionic detergents improve the analysis of membrane proteins by two-dimensional electrophoresis”

Electrophoresis **19**, 1901-1909.

Choi, J. K., Tak, K. H., Jin, L. T., Hwang, S. Y., Kwon, T. I. und Yoo, G. S. (2002)

“Background-free, fast protein staining in sodium dodecyl sulfate polyacrylamide gel using counterion dyes, zincon and ethyl violet”

Electrophoresis **23**, 4053-4059.

Chomczynski, P. und Sacchi, N. (1987)

“Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction.”

Anal. Biochem. **162**, 156-159.

Clary, D. O., Griff, I. C. und Rothman, J. E. (1990)

“SNAPs, a family of NSF attachment proteins involved in intracellular membrane fusion in animals and yeast”

Cell **61**, 709-721.

Corbett, J. M., Dunn, M. J., Posch, A. und Görg, A. (1994)

“Positional reproducibility of protein spots in two-dimensional polyacrylamide gel electrophoresis using immobilised pH gradient isoelectric focusing in the first dimension: an interlaboratory comparison”

Electrophoresis **15**, 1205-1211.

Cordon-Cardo, C., O'Brien, J. P., Casals, D., Rittman-Grauer, L., Biedler, J. L., Melamed, M. R. und Bertino, J. R. (1989)

“Multidrug-resistance gene (P-glycoprotein) is expressed by endothelial cells at blood-brain barrier sites”

Proc. Natl. Acad. Sci. U S A **86**, 695-698.

Dallas-Yang, Q., Jiang, G. und Sladek, F. M. (1998)

“Avoiding false positives in colony PCR”

BioTechniques **24**, 580-582.

D'Amore, P. A. (1990)

“Culture and study of pericytes”

in: Cell culture techniques in heart and vessel research (H. M. Piper, Hrsg.), 299-314, Springer-Verlag Berlin.

Danno, S., Nishiyama, H., Higashitsuji, H., Yokoi, H., Xue, J. H., Itoh, K., Matsuda, T. und Fujita, J. (1997)

“Increased transcript level of RBM3, a member of the glycine-rich RNA-binding protein family, in human cells in response to cold stress”

Biochem. Biophys. Res. Commun. **236**, 804-807.

De Moreno, M. R., Smith, J. F. und Smith, R.V. (1985)

“Silver staining of proteins in polyacrylamide gels: increased sensitivity through a combined Coomassie blue-silver stain procedure”

Anal. Biochem. **151**, 466-470.

Denis, M. (1986)

“Structure and function of cytochrome-c oxidase”

Biochimie **68**, 459-470.

Desilva, M. G., Lu, J., Donadel, G., Modi, W. S., Xie, H., Notkins, A.L. und Lan, M. S. (1996)

“Characterization and chromosomal localization of a new protein disulfide isomerase, PDIp, highly expressed in human pancreas”

DNA Cell. Biol. **15**, 9-16.

Eisenblätter, T. und Galla, H. J. (2002)

“A new multidrug resistance protein at the blood-brain barrier”

Biochem. Biophys. Res. Commun. **293**, 1273-1278.

El Hafny, B., Bourre, J. M. und Roux, F. (1996)

“Synergistic stimulation of  $\gamma$ -glutamyl transpeptidase and alkaline phosphatase activities by retinoic acid and astroglial factors in immortalized rat brain microvessel endothelial cells”

J. Cell Physiol. **167**, 451-460.

Feinberg, A. P. und Vogelstein, B. A. (1983)

“A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity”

Anal. Biochem. **132**, 6-13.

Fenn, J. B., Mann, M., Meng, C. K., Wong, S. F. und Whitehouse, C. M. (1989)

“Electrospray ionization for mass spectrometry of large biomolecules”

Science **246**, 64-71.

Fialka, I., Pasquali, C., Lottspeich, F., Ahorn, H. und Huber, L. A. (1997)

“Subcellular fractionation of polarized epithelial cells and identification of organelle-specific proteins by two-dimensional gel electrophoresis”

Electrophoresis **18**, 2582-2590.

Frey, A. (1991)

“Die  $\gamma$ -Glutamyltranspeptidase in der Blut-Hirn-Schranke des Schweines: Analyse der Primärstruktur und zelluläre Lokalisation”

Dissertation, TH-Darmstadt.

Ganong, B. R. und Delmore, J. P. (1991)

“Phase separation temperatures of mixtures of Triton X-114 and Triton X-45: application to protein separation”

Anal. Biochem. **193**, 35-37.

Gelsthorpe, M., Pulumati, M., McCallum, C., Dang-Vu K. und Tsubota, S. I. (1997)

“The putative cell cycle gene, enhancer of rudimentary, encodes a highly conserved protein found in plants and animals”

Gene **186**, 189-195.

Gevaert, K. und Vandekerckhove, J. (2000)

“Protein identification methods in proteomics”

Electrophoresis **21**, 1145-1154.

Gieffers, C., Koriath, F., Heimann, P., Ungermann, C. und Frey, J. (1997)

“Mitofilin is a transmembrane protein of the inner mitochondrial membrane expressed as two isoforms”

Exp. Cell. Res. **232**, 395-399.

Goldstein, G. W. und Betz., A. L. (1986)

“Die Blut-Hirn-Schranke”

Spektrum der Wissenschaft **11**, 82-91.

- Görg, A., Obermaier, C., Boguth, G., Csordas, A., Diaz, J. J. und Madjar, J. J. (1997)  
“Very alkaline immobilized pH gradients for two-dimensional electrophoresis of ribosomal and nuclear proteins”  
*Electrophoresis* **18**, 328-337.
- Gruhn, P. (2002)  
“Charakterisierung von spezifischen Blut-Hirn-Schranke-Proteinen aus Schwein”  
Diplomarbeit, TU-Darmstadt.
- Gygi, S. P., Rochon, Y., Franza, B. R., und Aebersold, R. (1999)  
“Correlation between protein and mRNA abundance in yeast”  
*Mol. Cell. Biol.* **19**, 1720-1730.
- Haebel, S., Albrecht, T., Sparbier, K., Walden, P., Korner, R. und Steup, M. (1998)  
“Electrophoresis-related protein modification: alkylation of carboxy residues revealed by mass spectrometry”  
*Electrophoresis* **19**, 679-686.
- Hendrick, J. P. und Hartl, F. U. (1993)  
“Molecular chaperone functions of heat-shock proteins”  
*Annu. Rev. Biochem.* **62**, 349-384.
- Henningsen, R., Gale, B. L., Straub, K. M. und De Nagel, D. C. (2002)  
“Application of zwitterionic detergents to the solubilisation of integral membrane proteins for two-dimensional gel electrophoresis and mass spectrometry”  
*Proteomics* **2**, 1479-1488.
- Herbert, B. (1999)  
“Advances in protein solubilisation for two-dimensional electrophoresis”  
*Electrophoresis* **20**, 660-663.
- Herbert, B. R., Molloy, M. P., Gooley, A. A., Walsh, B. J., Bryson, W. G. und Williams, K. (1998)  
“Improved protein solubility in two-dimensional electrophoresis using tributyl phosphine as reducing agent”  
*Electrophoresis* **19**, 845-851.

Himmel, H. M., Rasmusson, R. L. und Strauss, H. C. (1994)

“Agonist-induced changes of  $[Ca^{2+}]_i$  and membrane currents in single bovine aortic endothelial cells”  
Am. J. Physiol. **267**, C1338-C1350.

Hochstrasser, D. F., Harrington, M. G., Hochstrasser, A. C., Miller, M. J. und Merrill, C. R. (1988)

“Methods for increasing the resolution of two-dimensional protein electrophoresis”  
Anal. Biochem. **173**, 424-435.

Hoffman, G. G., Lee, S., Christiano, A. M., Chung-Honet, L. C., Cheng, W., Katchman, S., Uitto, J. und Greenspan, D. S. (1993)

“Complete coding sequence, intron/exon organization, and chromosomal location of the gene for the core I protein of human ubiquinol-cytochrome c reductase”  
J. Biol. Chem. **268**, 21113-21119.

Honore, B. (2000)

“The hnRNP 2H9 gene, which is involved in the splicing reaction, is a multiply spliced gene”  
Biochim. Biophys. Acta. **1492**, 108-119.

Huber, R., Romisch, J. und Paques, E. P. (1990)

“The crystal and molecular structure of human annexin V, an anticoagulant protein that binds to calcium and membranes”  
EMBO J. **9**, 3867-3874.

Hudmon, A. und Schulman, H. (2002)

“Neuronal  $Ca^{2+}$ /calmodulin-dependent protein kinase II: The Role of structure and autoregulation in cellular function”  
Annu. Rev. Biochem. **71**, 473-510.

Hunt, S. P. und Livesey, F. J. (2000)

“Functional genomics”  
Oxford University Press, Oxford.

Hutchison, H. T., Eisenberg, H. M. und Haber, B. (1985)

“High-affinity transport of glutamate in rat brain microvessels”  
Exp. Neurol. **87**, 260-269.



Ikegawa, S., Isomura, M., Koshizuka, Y. und Nakamura, Y. (1999)

“Cloning and characterization of a novel gene (C8orf2), a human representative of a novel gene family with homology to *C. elegans* C42.C1.9”

Cytogenet. Cell Genet. **85**, 227-237.

Jefferies, W. A., Brandon, M. R., Hunt, S. V., Williams, A. F., Gatter, K. C. und Mason, D.Y. (1984)

“Transferrin receptor on endothelium of brain capillaries”

Nature **312**, 162-163.

Kaplan, J. H. (2002)

“Biochemistry of Na,K-ATPase”

Annu. Rev. Biochem. **71**, 511-535.

Karas, M. und Hillenkamp, F. (1988)

“Laser desorption ionization of proteins with molecular masses exceeding 10,000 daltons”

Anal. Chem. **60**, 2299-2301.

Kim, J., Kim, S. H., Lee, S. U., Ha, G. H., Kang, D. G., Ha, N., Ahn, J. S., Cho, H. Y., Kang, S., Lee, Y., Hong, S. C., Ha, W. S., Bae, J. M., Lee, C. und Kim, J. W. (2002)

“Proteome analysis of human liver tumor tissue by two-dimensional gel electrophoresis and matrixassisted laser desorption/ionisation-mass spectrometry for identification of disease-related proteins”

Electrophoresis **23**, 4142-4156.

Kimelberg, H. K. und Norenberg, M. D. (1989)

“Astrocyten und Hirnfunktion”

Spektrum der Wissenschaft **6**, 52-60.

Klose, J. (1975)

“Protein mapping by combined isoelectric focusing and electrophoresis of mouse tissues. A novel approach to testing for induced point mutations in mammals”

Humangenetik **26**, 231-243.

Krause, D., Kunz, J. und Dermietzel, R. (1993)

“Cerebral pericytes - a second line of defense in controlling blood-brain barrier peptide metabolism”

Adv. Exp. Med. Biol. **331**, 149-152.

Laemmli, U. K. (1970)

“Cleavage of structural proteins during the assembly of the head of bacteriophage T4”

Nature **227**, 680-685.

Lahm, H. W. und Langen, H. (2000)

“Mass spectrometry: a tool for the identification of proteins separated by gels”

Electrophoresis **21**, 2105-2114.

Li, G., Laabich, A., Liu, O., Xue, J., und Cooper, N. (2000)

“Molecular cloning and sequence analyses of calcium/calmodulin-dependent protein kinase II from fetal and adult human brain”

Mol. Biol. Rep. **28**, 35-41.

Liang, P. und Pardee, A. B. (1992)

“Differential display of eukaryotic messenger RNA by means of the polymerase chain reaction”

Science **257**, 967-971.

Link, A. J., Eng, J., Schieltz, D. M., Carmack, E., Mize, G. J., Morris, D. R., Garvik, B. M. und Yates, J. R. (1999)

“Direct analysis of protein complexes using mass spectrometry”

Nature **17**, 676-682.

Loo, J. A. (1997)

“Studying noncovalent protein complexes by electrospray ionization mass spectrometry”

Mass. Spectrom. Rev. **16**, 1-23.

Lottspeich, F. und Zorbas., H. (Hrsg) (1998)

“Bioanalytik”

Spektrum Akademischer Verlag, Heidelberg Berlin.

Mandon, E. C., Ehses, I., Rother, J., van Echten, G. und Sandhoff, K. (1992)

“Subcellular localization and membrane topology of serine palmitoyltransferase, 3-dehydrosphinganine reductase, and sphinganine N-acyltransferase in mouse liver”

J. Biol. Chem. **267**, 11144-11148.

Mantych, G. J., James, D. E. und Devaskar, S. U. (1993)

“Jejunal/kidney glucose transporter isoform (glut-5) is expressed in human blood-brain barrier”

Endocrinology **132**, 35-40.

Meister, A. und Tate, S. S. (1976)

“Glutathione and related  $\gamma$ -Glutamyl compounds. Biosynthesis and utilization”

Annu. Rev. Biochem. **45**, 559-604.

Merril, C. R. (1990)

“Gel-staining techniques”

Methods Enzymol. **182**, 477-488.

Meyer, J., Mischeck, U., Veyhl, M., Henzel, K. und Galla, H. J. (1990)

“Blood-brain barrier characteristic enzymatic properties in cultured brain capillary endothelial cells”

Brain Res. **514**, 305-309.

Mischeck, U., Meyer, J. und Galla, H. J. (1989)

“Characterization of  $\gamma$ -Glutamyltranspeptidase activity of cultured endothelial cells from porcine brain capillaries”

Cell Tissue Res. **256**, 221-226.

Möckel, B., Zinke, H., Flach, R., Weiss, B., Weiler-Guttler, H. und Gassen, H. G. (1994)

“Expression of apolipoprotein A-I in porcine brain endothelium in vitro”

J. Neurochem. **62**, 788-798.

Molloy, M. P. (2000b)

“Two-dimensional electrophoresis of membrane proteins using immobilized pH gradients”

Anal. Biochem. **280**, 1-10.

Molloy, M. P., Herbert, B. R., Slade, M. B., Rabilloud, T., Nouwens, A. S., Williams, K. L. und Gooley, A. A. (2000a)

“Proteomic analysis of the Escherichia coli outer membrane”

Eur. J. Biochem. **267**, 2871-2881.

Molloy, M. P., Herbert, B. R., Walsh, B. J., Tyler, M. I., Traini, M., Sanchez, J. C., Hochstrasser, D. F., Williams, K. L. und Gooley, A. A. (1998)

“Extraction of membrane proteins by differential solubilisation for separation using two-dimensional gel electrophoresis”

Electrophoresis **19**, 867-844.

Mullis, K. B. (1987)

“Specific synthesis of DNA in vitro via a polymerase catalyzed chain reaction”

Methods Enzymol. **155**, 335-350.

Neuhoff, V., Arold, N., Taube, D. und Ehrhardt, W. (1988)

“Improved staining of proteins in polyacrylamide gels including isoelectric focusing gels with clear background at nanogram sensitivity using Coomassie Brilliant Blue G-250 and R-250”

Electrophoresis **9**, 255-262.

Neuhoff, V., Stamm, R., Pardowitz, I., Arold, N., Ehrhardt, W. und Taube, D. (1990)

“Essential problems in quantification of proteins following colloidal staining with coomassie brilliant blue dyes in polyacrylamide gels, and their solution”

Electrophoresis **11**, 101-117.

Nijtmans, L., Artal, S., Grivell, L. und Coates, P. (2002)

“The mitochondrial PHB complex: roles in mitochondrial respiratory complex assembly, ageing and degenerative disease”

Cell. Mol. Life Sci. **59**, 143-155.

Odgren, P. R., Toukatly, G., Bangs, P. L., Gilmore, R. und Fey, E.G. (1996)

“Molecular characterization of mitofilin (HMP), a mitochondria-associated protein with predicted coiled coil and intermembrane space targeting domains”

J. Cell Sci. **109**, 2253-2264.

O'Farrell, P. H. (1975)

“High resolution two-dimensional electrophoresis of proteins”

J. Biol. Chem. **250**, 4007-4021.

O'Farrell, P. Z., Goodman, H. M. und O'Farrell, P. H. (1977)

“High resolution two-dimensional electrophoresis of basic as well as acidic proteins”

Cell **12**, 1133-1141.

Owczarek, C. M., Treutlein, H. R., Portbury, K. J., Gulluyan, L. M., Kola, I. und Hertzog, P. J. (2001)  
“A novel member of the STOMTIN/EPB72/mec-2 family, stomatin-like 2 (STOML2), is ubiquitously expressed and localizes to HSA chromosome 9p13.1”

Cytogenet. Cell Genet. **92**, 196-203.

Perrin, B. J. und Huttenlocher., A. (2002)

“Calpain”

Int. J. Biochem. Cell Biol. **34**, 722-725.

Pinkstaff, J., Chappell, S., Mauro, V., Edelman, G. und Krushel, L. (2001)

“Internal initiation of translation of five dendritically localized neuronal mRNAs”

Proc. Natl. Acad. Sci. USA **98**, 2770-2775.

Poehling, H. und Neuhoff, V. (1981)

“Visualization of proteins with a silver stain: a critical analysis”

Electrophoresis **2**, 141-147.

Pogge von Strandmann, E., Senkel, S. und Ryffel, G. U. (2001)

“ERH (enhancer of rudimentary homologue), a conserved factor identical between frog and human, is a transcriptional repressor”

Biol. Chem. **382**, 1379-1385.

Rabilloud, T. (1998a)

“Use of thiourea to increase the solubility of membrane proteins in two dimensional electrophoresis with immobilized pH gradients”

Electrophoresis **19**, 758-760.

Rabilloud, T., Adessi, C., Giraudel, A. und Lunardi, J. (1997)

“Improvement of the solubilization of proteins in two-dimensional electrophoresis with immobilized pH gradients”

Electrophoresis **18**, 307-316.

Rabilloud, T., Gianazza, E., Catto, N. und Righetti, P. G. (1990)

“Amidosulfobetaines, a family of detergents with improved solubilization properties: application for isoelectric focusing under denaturing conditions”

Anal. Biochem. **185**, 94-102.

Rabilloud, T., Kieffer, S., Procaccio, V., Louwagie, M., Courchesne, P. L., Patterson, S. D., Martinez, P., Garin, J. und Lunardi, J. (1998b)

“Two-dimensional electrophoresis of human placental mitochondria and protein identification by mass spectrometry: toward a human mitochondrial proteome”

Electrophoresis **19**, 1106-1014.

Ramsby, M. L. und Makowski, G. S. (1999)

“Differential detergent fractionation of eucariotic cells. Analysis by two-dimensional gel Electrophoresis”

Methods Mol. Biol. **112**, 53-66.

Rao, R. V., Peel, A., Logvinova, A., del Rio, G., Hermel, E., Yokota, T., Goldsmith, P. C., Ellerby, L. M., Ellerby, H.M. und Bredesen, D. E. (2002)

“Coupling endoplasmic reticulum stress to the cell death program: role of the ER chaperone GRP78”

FEBS Lett. **514**, 122-128.

Righetti, P. G., Tudor, G. und Gianazza, E. (1982)

“Effect of 2-mercaptoethanol on pH gradients in isoelectric focusing”

J. Biochem. Biophys. Methods **6**, 219-227.

Risau, W., Hallermann, R. und Albrecht, U. (1986)

“Differentiation-dependent expression of Proteins in brain endothelium during development of the blood-brain barrier”

Dev. Biol. **117**, 537-545.

Rouquie, D., Peltier, J. B., Marquis-Mansion, M., Tournaire, C., Dumas, P. und Rossignol, M. (1997)

“Construction of a directory of tobacco plasma membrane proteins by combined two-dimensional gel electrophoresis”

Electrophoresis **18**, 654-660.

Sambrook, J., Fritsch, E. F. und Maniatis, T. (1989)

“Molecular cloning - al laboratory manual”

Cold Spring Laboratory Press, New York.

Sanchez del Pino, M. M., Peterson, D. R. und Hawkins, R. A. (1995)

“Neutral amino acid transport characterization of isolated luminal and abluminal membranes of the blood-brain barrier”

J. Biol. Chem. **270**, 14913-14918.

Sanger, F., Micklen, S. und Coulson, A. R. (1977)

“DNA-sequencing with chain-terminating inhibitors”

Proc. Natl. Acad. Sci. USA **74**, 5463-5467.

Santoni, V., Rabilloud, T., Doumas, P., Rouquie, D., Mansion, M., Kieffer, S., Garin, J. und Rossignol, M. (1999)

“Towards the recovery of hydrophobic proteins on two-dimensional electrophoresis gels”

Electrophoresis **20**, 705-711.

Scheetz, A., Nairn, A. und Constantine-Paton, M (2000)

“NMDA receptor-mediated control of protein synthesis at developing synapses”

Nat. Neurosci **3**, 211 - 216.

Scheler, C., Lamer, S., Pan, Z., Li, X. P., Salnikow, J. und Jungblut, P. (1998)

“Peptide mass fingerprint sequence coverage from differently stained proteins on two-dimensional electrophoresis patterns by matrix assisted laser desorption/ionization-mass spectrometry (MALDI-MS)”

Electrophoresis **19**, 918-927.

Schroeter, M. L., Mertsch, K., Giese, H., Muller, S., Sporbert, A., Hickel, B. und Blasig, I. E. (1999)

“Astrocytes enhance radical defence in capillary endothelial cells constituting the blood-brain barrier”

FEBS Lett. **449**, 241-244.

Shi, Y., Zhai, H., Wang, X., Wu, H., Ning, X., Han, Y., Zhang, D., Xiao, B., Wu, K. und Fan, D. (2002)

“Multidrug-resistance-associated protein MGr1-Ag is identical to the human 37-kDa laminin receptor precursor”

Cell. Mol. Life Sci. **59**, 1577-1583.

Sickmann, A., Dormeyer, W., Wortelkamp, S., Voitalla, D., Kuhn, W. und Meyer, H. (2000)  
“Identification of proteins from human cerebrospinal fluid, separated by two-dimensional polyacrylamide gel electrophoresis”  
Electrophoresis **21**, 2721-2728.

Sims, D. E. (1986)  
“The pericyte - a review”  
Tissue Cell **18**, 153-174.

Smith, L. M., Sanders, J. Z., Kaiser, R. J., Hughes, P. Dodd, C., Conell, C. R., Heiner, C., Kent, S. B. und Hood, L. E. (1986)  
“Fluorescence detection in automated DNA sequence analysis”  
Nature **321**, 674-679.

Steel, G. J., Buchheim, G., Edwardson, J. M. und Woodman, P. G. (1997)  
“Evidence for interaction of the fusion protein alpha-SNAP with membrane lipid”  
Biochem. J. **325**, 511-518.

Stoll, J., Wadhvani, K. C. und Smith, Q. R. (1993)  
“Identification of the cationic amino acid transporter (System y<sup>+</sup>) of the rat blood-brain barrier”  
J. Neurochem. **60**, 1956-1959.

Sudo, K., Takahashi, E. und Nakamura, Y. (1995)  
“Isolation and mapping of the human EIF4A2 gene homologous to the murine protein synthesis initiation factor 4A-II gene Eif4a2”  
Cytogenet. Cell Genet. **71**, 358-388.

Tayarani, I., Lefauconnier, J. M., Roux, F. und Bourre, J. M. (1987)  
“Evidence for an alanine, serine, and cysteine system of transport in isolated brain capillaries”  
J. Cereb. Blood Flow Metab. **7**, 585-591.

Teifel, M. (1996)  
“Transfektion vaskulärer Endothelzellen. Optimierung der Transfektion und Anwendung zur heterologen Genexpression”  
Dissertation, TH-Darmstadt.



- Thongboonkerd, V., Mc Leish, K. R., Arthur, J. M. und Klein, J. B. (2002)  
"Proteomic analysis of normal human urinary proteins isolated by acetone precipitation or ultracentrifugation"  
Kidney Int. **62**, 1461-1469.
- Tobimatsu, T. und Fujisawa, H. (1989)  
"Tissue-specific expression of four types of rat calmodulin-dependent protein kinase II mRNAs"  
J. Biol. Chem. **264**, 17907-17912.
- Tsukakara, F., Yoshioka, T. und Muraki, T. (2000)  
"Molecular and functional characterization of HSC54, a novel variant of human heat-shock cognate protein 70"  
Mol. Pharmacol. **58**, 1257-1263.
- Vachon, V., Pouliot, J. F., Laprade, R. und Beliveau, R. (1991)  
"Fractionation of renal brush border membrane proteins with Triton X-114 phase partitioning"  
Biochem. Cell Biol. **69**, 206-211.
- Vergeres, W. und Waskell, L. (1995)  
"Cytochrome b5, its functions, structure and membrane topology"  
Biochimie **77**, 604-620.
- Vito, P., Lacana, E. und D'Adamio, L. (1996)  
"Interfering with apoptosis: Ca(2+)-binding protein ALG-2 and Alzheimer's disease gene ALG-3"  
Science **271**, 521-525.
- Wakui, S., Furusato, M., Muto, T., Ohshige, H., Takahashi, H. und Ushigome, S. (1997)  
"Transforming growth factor-beta and urokinase plasminogen activator presents at endothelial cell-pericyte interdigitation in human granulation tissue"  
Microvasc Res **54**, 262-269.
- Wang, Y. und Morrow, J. S. (2000)  
"Identification and characterization of human SLP-2, a novel homologue of stomatin (band 7.2b) present in erythrocytes and other tissues"  
J. Biol. Chem. **275**, 8062-8071.

Washburn, M. P., Wolters, D. und Yates, J. R. (2001)

“Large-scale analysis of the yeast proteome by multidimensional protein identification technology”  
Nat. Biotechnol. **19**, 242-247.

Weiler-Güttler, H., Zinke, H., Möckel, B., Frey, A. und Gassen, H. G. (1989)

“cDNA cloning and sequence analysis of glucose transporter from porcine blood-brain barrier”  
Biol. Chem. Hoppe Seyler **370**, 467-473.

Weiss, B. und Stoffel, W. (1997)

“Human and murine serine-palmitoyl-CoA transferase - cloning, expression and characterization of the key enzyme in sphingolipid synthesis”  
Eur. J. Biochem. **249**, 239-247.

Wen, L., Huang, J. K., Johnson, B. H. und Reeck, G. R. (1989)

“A human placental cDNA clone that encodes nonhistone chromosomal protein HMG-1”  
Nucleic Acids Res. **17**, 1197-1214.

Westermeier, R., Postel, W., Weser, J. und Görg, A. (1983)

“High-resolution two-dimensional electrophoresis with isoelectric focusing in immobilized pH gradients”  
J. Biochem. Biophys. Methods **8**, 321-330.

Wible, B. A., De Biasi, M., Majumder, K., Tagliatela, M. und Brown, A. M. (1995)

“Cloning and functional expression of an inwardly rectifying K<sup>+</sup> channel from human atrium”  
Circ. Res. **76**, 343-350.

Wissing, J., Heim, S., Flohe, L., Bilitewski, U. und Frank, R. (2000)

“Enrichment of hydrophobic proteins via Triton X-114 phase partitioning and hydroxyapatite column chromatography for mass spectrometry”  
Electrophoresis **21**, 2589-2593.

Wolburg, H., Neuhaus, J., Kniesel, U., Krauss, B., Schmid, E. M., Ocalan, M., Farrell, C. und Risau, W. (1994)

“Modulation of tight junction structure in blood-brain barrier endothelial cells. Effects of tissue culture, second messengers and cocultured astrocytes”  
J. Cell Sci. **107**, 1347-1357.

Yoo, M. (1997)

“Identification of two homologous cytochrome b5s in rat brain”

Biochem. Biophys. Res. Commun. **236**, 641-642.

Zheng, X. M., Black, D., Chambon, P. und Egly, J. M. (1990)

“Sequencing and expression of complementary DNA for the general transcription factor BTF3”

Nature **344**, 556-559.